
The background of the slide is an aerial photograph of the Arctic region, showing a vast expanse of broken sea ice. The ice floes are irregular in shape and size, ranging from small, thin pieces to larger, more substantial chunks. The water between the ice floes is a deep, dark blue, creating a complex, mosaic-like pattern. The overall color palette is dominated by various shades of blue, from light, almost white, to deep, dark navy blue.

# The Causes and Consequences of a Rapidly Changing Arctic

Patrick C. Taylor  
Climate Science Branch  
NASA Langley Research Center  
OneNOAA Seminar Series  
August 16, 2018



Source: cnn.com

GHERIEL BLOUIN/REUTERS/GETTY IMAGES  
Land  
edge  
onal  
erve

# What is the National Climate Assessment? What is the Climate Science Special Report?

- As part of its mandate, USGCRP is charged with leading a quadrennial assessment of U.S. climate: The National Climate Assessment.
- In an effort to provide a sustained assessment, the 4<sup>th</sup> National Climate Assessment will be released in two volumes.
- The CSSR is an authoritative assessment of the science of climate change in the U.S.

## CSSR Stats:

63 Key Findings

3 Coordinating Lead Authors

32 Lead Authors


19 Contributing Authors

470 pages

6 Reviews

3 Review Editors

2 Leaked copies



U.S. Global Change  
Research Program

<http://science2017.globalchange.gov>

# Arctic Changes and their Effect on Alaska and the Rest of the United States

The Climate Science Special Report  
U.S. Global Change Research Program

Chapter Lead: Patrick Taylor (NASA)

Lead Authors: J. Perlwitz (NOAA), D. J. Wuebbles (Univ. Illinois),  
W. Maslowski (NRL)

Contributing author: J. Walsh (Univ. of Alaska)

<https://science2017.globalchange.gov>



CSSR, Ch. 11, Key Finding 2: Rising Alaskan permafrost temperatures are causing permafrost to thaw and become more discontinuous; this process releases additional carbon dioxide and methane, resulting in an amplifying feedback and additional warming (*high confidence*). The overall magnitude of the permafrost–carbon feedback is uncertain; however, it is clear that these emissions have the potential to compromise the ability to limit global temperature increases.

Trends in permafrost temperatures range from 0.2 to 0.7°C per decade.




Figure 11.5

Methane Bubbles

Source: NASA ABoVE

September


1984

Sea Ice Age

0-1 1-2 2-3 3-4 4+  
Years


The character of Arctic sea ice has changed dramatically.

[View animation](#)



CSSR, Ch. 11 Key Finding 3:  
Arctic land and sea ice loss observed in the last three decades continues, in some cases accelerating (*very high confidence*)....Arctic-wide ice loss is expected to continue through the 21st century, very likely resulting in nearly sea ice-free late summers by the 2040s (*very high confidence*).

## Trends in Sea Ice Melt Season





Arctic sea ice melt season has lengthened since 1979.

Along the Alaskan northern and western coast, the season has lengthened by 20-30 days per decade adding up to 90 more sea ice free days annually.

# GRACE Observations of Greenland Ice Mass Changes

[View animation](#)



Dramatic changes have occurred across the Greenland Ice Sheet, particularly at its margins.

1941



2004




Figure 11.4

Muir Glacier has retreated more the 4 miles to the northwest.

CSSR, Ch. 11 Key Finding 4:

It is very likely that human activities have contributed to observed arctic surface temperature warming, sea ice loss, glacier mass loss, and Northern Hemisphere snow extent decline (*high confidence*).

CSSR, Ch. 11 Key Finding 5:


Atmospheric circulation patterns connect the climates of the Arctic and the contiguous United States...the midlatitude circulation has influenced observed arctic temperatures and sea ice (*high confidence*). However, confidence is low regarding whether or by what mechanisms observed arctic warming may have influenced the midlatitude circulation and weather patterns over the continental United States....



Source: NASA IceBridge


# Nature of Arctic Amplification

- Arctic Amplification Defined: Arctic warms more given the same CO<sub>2</sub> forcing
- Arctic warming focused at the surface.
- Arctic warming strongest in fall and winter.




# Processes influencing Arctic Amplification

Warm, Sun-lit season



Cold, dark season



Cohen et al. (in review)


[View animation](#)



[View animation](#)



# Hours above Freezing at Morris Jesup, Greenland




61 hours 2018

GFS/CFSR 2m T Anomaly (°C) [1979-2000 base]  
Init 2018/02/23 00Z | f000 Valid Fri 00Z, Feb 23, 2018

ClimateReanalyzer.org  
University of Maine | Climate Change Institute

View animation



- World + 0.6 °C
- NH + 1.2 °C
- SH + 0.1 °C
- Arctic + 5.9 °C
- Antarctic + 0.7 °C
- Tropics + 0.5 °C

 **Robert Rohde**  
@rarohde

Replying to @rarohde  
In 2018, there have already been 61 hours above freezing at Cape Morris Jesup, Greenland.


The previous record was 16 hours before the end of April in 2011.

8:02 PM - Feb 25, 2018

♡ 157 💬 180 people are talking about this


# A complete picture of Arctic Amplification?

LWDN is the dominant term contributed to Arctic Amplification and integrates the local and remote mechanisms.




# Are the interactions between local and remote mechanisms most important?


Local Mechanisms Only



Remote Mechanisms Only




Local+Remote Mechanisms



# 2016-17 Arctic Sea Ice Growth Season



2016-17 also exhibited low sea ice extent from October-March, well below recent averages, contributing to one of the lowest end of season Arctic sea ice volume on record.



Hegyi and Taylor (2018; GRL): The Unprecedented 2016–2017 Arctic Sea Ice Growth Season: The Crucial Role of Atmospheric Rivers and Longwave Fluxes

November 17, 2016



Presence of a moisture intrusion



Increased downwelling LW fluxes


November 17, 2016  
 a) Anomalous PW (shading), 925 hPa winds (vectors), and area of moisture intrusion (red contour)  
 b) Anomalous downwelling LW flux anomaly  
 c) Sea ice volume growth  
 d) Sea ice extent retreat

Sea ice volume loss



Sea ice extent retreat


# Arctic Temperature Response (RCP 8.5)



Most warming in fall/winter



More model disagreement in the Arctic than any other region

Bottom heavy warming profile



# Projected Arctic Surface Temperature Warming (RCP8.5)

Annual Mean      Winter (JF)      Sunlit (MAMJJAS)      Autumn (OND)



Largest differences between CMIP5 models occur in fall and winter in the Barents-Kara Seas and the Chukchi-Beaufort Seas regions.





# Projected Arctic Sea Ice Loss (RCP8.5)

Annual Mean





Winter (JF)

Sunlit (MAMJJAS)

Autumn (OND)



Ensemble Mean RCP8.5  $\Delta$ SIC by 2100 (%)




Model Ensemble Std. Dev. in  $\Delta$ SIC

3 6 9 12 15 18 21 24 27 30 33 36 39


Largest differences between CMIP5 models occur in fall and winter in the Barents-Kara Seas and the Chukchi-Beaufort Seas regions.

# Projected Arctic Sea Ice Loss (RCP4.5 vs. RCP8.5)


## September Arctic Sea Ice Projections





1985-2005 Average  
February September



2081-2100 Average,  
February RCP4.5 September



2081-2100 Average,  
February RCP8.5 September



In recent decades, Alaskan wildfire activity has increased in both boreal forest and in the Arctic tundra.

Image credit: NASA  
MODIS (showing 314  
active fires on June 25,  
2015)

- Large wildfires ( $>100,000 \text{ km}^2$ ) have *likely* become more frequent since 1959.
- Human activities have *likely* lengthened the wildfire season and increased the risk of severe fires.

A plethora of opportunities and challenges  
in a changing Arctic...



Source: NASA IceBridge

Thank you.

patrick.c.taylor@nasa.gov

<https://science2017.globalchange.gov/chapter/11/>

Questions?

Source: NASA IceBridge